

RESEARCH INSTITUTE OF APPLIED PHYSICS
TASHKENT STATE UNIVERSITY

Preprint NIIPF-93/07

QUANTUM DEFORMATIONS FOR THE DIAGONAL R-MATRICES
Talk at Workshop on Supersymmetry and Quantum Groups, Dubna, 15-20
July 1993

B.M.ZUPNIK

Tashkent - 1993

QUANTUM DEFORMATIONS FOR THE DIAGONAL R-MATRICES

B.M.ZUPNIK

Research Institute of Applied Physics, Tashkent State University, Vuzgorodok, Tashkent
700095, Uzbekistan

Abstract

We consider two different types of deformations for the linear group $GL(n)$ which correspond to using of a general diagonal R-matrix. Relations between braided and quantum deformed algebras and their coactions on a quantum plane are discussed. We show that tensor-grading-preserving differential calculi can be constructed on braided groups, quantum groups and quantum planes for the case of the diagonal R-matrix.

Deformations of the linear group $GL(n)$, vector spaces and commutative algebras are intensively discussed in the theory of quantum groups [1-4]. More simple (G, f) -graded deformations of Lie algebras are well known [5,6] and recently their R-matrix generalizations (braided algebras) have been considered [7,8].

We shall discuss the simplest multiparametric deformations of the commutative algebra and linear group generating by a diagonal R_D -matrix [9]. The general diagonal unitary R-matrix has the following form:

$$(R_D)_{km}^{ij} = q(ji)\delta_k^i\delta_m^j \quad (1)$$

where $q(ji)$ are complex deformation parameters

$$q(ji) = 1/q(ij), \quad q(ii) = 1 \quad (2)$$

This R-matrix is a partial case $X = 1$ of the multiparametric nondiagonal R-matrix [10].

Let us define a quantum plane [2] as the formal-series algebra with generators x^i ($i = 1, 2, \dots, n$)

$$x^i x^j = (R_D)_{km}^{ji} x^k x^m = q(ij)x^j x^i \quad (3)$$

One can treat the deformed groups as different cotransformations on the quantum plane. We shall study two types of $GL(n)$ deformations using R_D -matrix (1): braided group $DGL(n)$ and quantum group $GL_D(n)$. Symbol D in our notations corresponds to the diagonal structure of R_D -matrix.

One can use the natural tensor I_D -grading on the quantum plane and corresponding differential calculus. Here I_D is a commutative semigroup, which can be defined as the set of covariant and contravariant multiindices with the multiplication rule *

$$I\left(\frac{p}{r}\right) = \left(\frac{i_1 \dots i_p}{k_1 \dots k_r}\right), \quad I\left(\frac{p}{r}\right) * I\left(\frac{s}{t}\right) = I\left(\frac{p+s}{r+t}\right) \quad (4)$$

The order of upper or low indices is unessential for the tensor grading so we can use some type of symmetrization of indices, for instance, D-symmetrization. Note, that we must distinguish the multiindices with equal number indices of different sort, e.g. $(123) \neq (345)$. Nevertheless one can consider the equivalence relation E in the set I_D : **the multiindices are equivalent if they have some identical subset of indices and also an arbitrary number of upper and low indices coinciding in pairs**, for example

$$\left(\frac{12}{34}\right) \sim \left(\frac{121}{341}\right) \sim \left(\frac{1235}{3435}\right) \sim \left(\frac{12357}{34357}\right) \quad (5)$$

Let us denote $\tilde{I}_D = I_D/E$ the corresponding factorset which has also a multiplication rule for classes of multiindices. Consider the generators dx^i , ∂_i of the special differential calculus preserving $\tilde{I}_D$ -grading [9]

$$\begin{aligned} dx^i x^k &= [ik] x^k dx^i \\ dx^i dx^k &= -[ik] dx^k dx^i \\ \partial_i x^k &= \delta_i^k + [ki] x^k \partial_i \\ \partial_i \partial_k &= [ik] \partial_k \partial_i \\ \partial_k dx^i &= [ik] dx^i \partial_k \end{aligned} \quad (6)$$

Here the notation $[ik] = q(ik)$ is used.

Let us introduce the following notations for $\tilde{I}_D$ -graded algebras: $M_D(n) = M_D(x)$ is the algebra with generators x^i , $\Lambda_D(n)$ is the external algebra with generators x^i, dx^i and $D_1(M_D)$ is a tangent vector space of first-order differential operators on $M_D(n)$. $D_1(M_D)$ can be treated as the infinite-dimensional Lie D-algebra with the generalized D-commutator of basic elements [9]

$$D_k^{I(p)} = x^{I(p)} \partial_k = x^{i_1} \dots x^{i_p} \partial_k \quad (7)$$

Consider a simple braided D-deformations of the linear group $DGL(n)$ and corresponding Lie algebra $dgl(n)$ [5,6,9]. $DGL(n)$ is an unital algebra with noncommuting generators L_k^i

$$L_k^i L_m^j = [ij][jk][km][mi] L_m^j L_k^i = \left[\frac{ij}{k|m}\right] L_m^j L_k^i \quad (8)$$

where a special notation for the commutation factor is used. $M_D(n)$ and $DGL(n)$ are examples of D-commutative algebras. A structure of Hopf D-algebra on $DGL(n)$ is consistent with $\tilde{I}_D$ -grading

$$\begin{aligned} \Delta(L_k^i) &= L_j^i \otimes_q L_k^j = [ij][jk][ki] L_k^j \otimes_q L_k^i j \\ s(L_k^i) &= (L^{-1})_k^i, \quad \varepsilon(L_k^i) = \delta_k^i \end{aligned} \quad (9)$$

where a special D-symmetrical tensor product is used [9]. We can choose the exponential parametrization for the braided matrices

$$L_k^i = (\exp \lambda)_k^i = \delta_k^i + \lambda_k^i + \dots \quad (10)$$

$$\lambda_k^i = \lambda_m^j (M_j^m)_k^i, \quad (M_j^m)_k^i = \delta_j^i \delta_k^m \quad (11)$$

Here M_j^m are generators of the fundamental representation of $dgl(n)$ and $\lambda_m^j (M_j^m)$ is an element of the corresponding noncommutative envelope [9]. Note, that the matrix elements of $dgl(n)$ -generators are not deformed.

Cotransformation of $DGL(n)$ on the quantum plane $M_D(n)$ conserves $\tilde{I}_D$ -grading

$$x'^i = L_k^i \otimes_q x^k = [ik] x^k \otimes_q L_k^i \quad (12)$$

$$dx'^i = L_k^i \otimes_q dx^k \quad (13)$$

$$\partial'_i = \partial_k \otimes_q (L^{-1})_i^k \quad (14)$$

It is convenient to use the equivalence relations for the grading of matrices

$$L_k^i \sim (L^2)_k^i \sim (L^{-1})_k^i \sim (\ln L)_k^i \quad (15)$$

One can impose an additional restriction on the deformation parameters $[ik] = q(ik)$, which results in disappearance of deformations for $dgl(n)$

$$[ik][kj] = [ij] \implies [\frac{ij}{k|m}] = 1 \quad (16)$$

Note, that $dgl(2) \simeq gl(2)$ without any additional restriction. Using the Eq(16) one can define the coaction of commutative matrices L_k^i on the noncommutative space $M_D(n)$

$$\begin{aligned} L_k^i L_m^j &= L_m^j L_k^i \\ L_k^i \otimes_q x^l &= [il][lk] x^l \otimes_q L_k^i \end{aligned} \quad (17)$$

Now we shall consider a quantum deformation $GL_D(n)$ of the linear group for the diagonal R-matrix (1). $F(GL_D(n))$ is a Hopf algebra with generators a_k^i

$$R_D A_1 A_2 = A_2 A_1 R_D \implies a_k^i a_m^j = [ij][km] a_m^j a_k^i \quad (18)$$

Stress, that multiplication and comultiplication in $F(GL_D(n))$ conserve some tensor grading, but an antipode map change tensor grading

$$\begin{aligned} s(a_k^i) &= (a^{-1})_k^i \\ (a^{-1})_k^i (a^{-1})_m^j &= [ji][km] (a^{-1})_m^j (a^{-1})_k^i \\ a_k^i (a^{-1})_m^j &= [mi][kj] (a^{-1})_m^j a_k^i \end{aligned} \quad (19)$$

The $\tilde{I}_D$ -differential calculus on $M_D(n)$ (6) is covariant with respect to the coaction of the quantum group $GL_D(n)$

$$\begin{aligned} x'^i &= a_k^i \otimes x^k, & dx'^i &= a_k^i \otimes dx^k \\ \partial'_i &= \partial_k \otimes (a^{-1})_i^k = (a^{-1})_i^k \otimes \partial_k \end{aligned} \quad (20)$$

where the symmetrical tensor product is used.

Stress , that different deformations of the linear group are closely connected as different cotransformations of the quantum plane . Let us discuss the manifestations of this connection and examples of the combined using of $GL_D(n)$ and $DGL(n)$ structures :

A) Consider a differential calculus on the quantum group $GL_D(n)$

$$\begin{aligned} da_k^i a_m^j &= [ij][mk] a_m^j da_k^i \\ da_k^i da_m^j &= -[ij][mk] da_m^j da_k^i \\ da_k^i (a^{-1})_m^j &= [mi][kj] (a^{-1})_m^j da_k^i \end{aligned} \quad (21)$$

Right-invariant differential forms ρ_k^i on $GL_D(n)$ [3] can be treated as 1-forms with $dgl(n)$ -structure

$$\begin{aligned} \rho_k^i &= da_l^i (a^{-1})_k^l \\ \rho_k^i \rho_m^j &= - \left[\frac{ij}{k|m} \right] \rho_m^j \rho_k^i \end{aligned} \quad (22)$$

B) $DGL(n)$ -matrices generate "left orbit" on the quantum $GL_D(n)$ -matrices

$$\begin{aligned} a_k^i &\rightarrow \tilde{a}_k^i = L_j^i a_k^j \\ L_l^i a_s^r &= [ir][rl] a_s^r L_l^i \\ \tilde{a}_k^i \tilde{a}_m^j &= [ij][mk] \tilde{a}_m^j \tilde{a}_k^i \\ \tilde{a}_k^i a_m^j &= [ij][mk] a_m^j \tilde{a}_k^i \\ L_k^i &= \tilde{a}_j^i (a^{-1})_k^j \end{aligned} \quad (23)$$

Note , that $DGL(n)$ -matrices can be constructed in terms of quantum matrices.

C) Quantum GL_D -matrices can define "two-sided quantum orbit" on $DGL(n)$

$$L_k^i \leftarrow a_j^i L_m^j (a^{-1})_k^m \quad (24)$$

where the elements $a_j^i, (a^{-1})_k^m$ commute with L_m^i . This construction was used in Ref[11] for deformation of gauge fields .

Deformations of the special linear group $SL(n)$ for the diagonal R_D -matrix can be obtained by a factorization of quantum $GL(n)$ -matrices [12,13]

$$a_k^i = \hat{a}_k^i (D_q)^{1/n} \quad (25)$$

where D_q is the quantum determinant and $\hat{a}_k^i$ are generators of $SL_D(n)$ -group . Note , that a quantum deformation vanishes for $SL_D(2)$ group , however we have nontrivial one-parametric deformation for the case of $SL_D(3)$ quantum group.

Let $q(12), q(23), q(31)$ are independent deformation parameters of $GL_D(3)$ (18). Consider a quantum determinant of $GL_D(3)$ -matrix a_k^i

$$\begin{aligned} D_q &= a_1^1 a_2^2 a_3^3 + [21][31] a_2^1 a_3^2 a_1^3 + [31][32] a_3^1 a_2^3 a_1^2 - \\ &\quad [21] a_2^1 a_1^2 a_3^3 - [32] a_1^1 a_3^2 a_2^3 - [21][31][32] a_3^1 a_2^2 a_1^3 \end{aligned} \quad (26)$$

The quantum determinant is not central element in $GL_D(3)$. Eq(25) determines elements of the $SL_D(n)$ -matrix $\hat{a}_k^i$ which satisfy the commutation relations (18) with the following deformation parameter

$$\hat{q}(12) = \hat{q}(23) = \hat{q}(31) = ([12][23][31])^{1/3} \quad (27)$$

References

- [1] Reshetikhin N.Yu., Takhtajan L.A., Faddeev L.D. *Algeb. anal.* 1 (1989) 178.
- [2] Manin Yu.I. *Comm. Math. Phys.* 123 (1989) 163 ; *Teor. Mat. Phys.* 92 (1992) 425.
- [3] Woronowicz S.L. *Comm. Math. Phys.* 122 (1989) 125.
- [4] J.Wess , Zumino B. *Nucl. Phys.B (Proc. Suppl.)* 18 (1990) 302.
- [5] Scheunert M. *Jour. Math. Phys.* 20 (1979) 712.
- [6] Mosolova M.V. *Matem. zapis.* 29 (1981) 35.
- [7] Majid S. *Jour. Math. Phys.* 32 (1991) 3246.
- [8] Gurevich D.I. *Doklady AN SSSR*, 288 (1986) 797.
- [9] Zupnik B.M. *Teor. Mat. Fiz.* 95 (1993); Preprint NIIPF-92-01, HEP TH/9211065.
- [10] Schirmacher A. *Zeit. Phys. C50* (1991) 321.
- [11] Isaev A.P., Popowicz Z. Preprint JINR E2-93-54.
- [12] Ogievetsky O., Wess J. *Zeit. Phys. C50* (1991) 123.
- [13] Schupp P., Watts P., Zumino B. *Lett. Math. Phys.* 25 (1991) 139.